

Soil biodiversity

Paul Henning Krogh¹, Henning Petersen², Arwyn Jones³,
Luca Montanarella³, Ciro Gardi³ and Simon Jeffery³

¹Aarhus University, National Environmental Research Institute, Dpt. of Terrestrial Ecology

²Mols Laboratory, Natural History Museum, Aarhus

³European Commission, Joint Research Centre, Institute of Environmental Sustainability




NATIONAL ENVIRONMENTAL
RESEARCH INSTITUTE
AARHUS UNIVERSITY

NATURHISTORISK
MUSEUM
AARHUS

ies
Institute for
Environment and
Sustainability

The novel **EUROPEAN ATLAS OF SOIL BIODIVERSITY**, published by the Joint Research Centre, presents an overview of soil biodiversity in Europe and assesses threats and the contributions to ecosystem services. It is intended for raising awareness of soil biodiversity and its crucial role in sustaining fertility and as a driver of organic matter decomposition and carbon sequestration. The awareness is needed for the proposed Soil Framework Directive by the European Commission in an attempt to prevent further soil degradation across the European Union, and to repair the damage that has already been done.




The awareness is needed for the proposed Soil Framework Directive by the European Commission in an attempt to prevent further soil degradation across the European Union, and to repair the damage that has already been done.

A Danish beech forest is inhabited by about 1000 soil invertebrate species

Diversity was assessed in the Hestehave beech forest project, a main Danish contribution to IBP (International Biological Programme: 1964-1974). The project included an extensive soil biological study of soil microflora and the most important soil fauna groups. The pie diagram illustrates the species richness found in 3 hectares of beech forest soil based on the Hestehave results supplemented with results from similar German beech forests.

Earthworm biomass in grassland soil is equivalent to having up to 5 cows per hectare

A normal grassland field can sustain 1 cow per hectare. The biomass of all soil organisms amounts to 10 tons per hectare.



Estimated number of species among almost all taxonomic invertebrate groups found in a Danish beech forest floor.

Approximate range of biomass of each major component of the biota in a typical temperate grassland soil.

| | Biomass t ha ⁻¹ | Cow equivalents |
|--------------------|----------------------------|-----------------|
| Plant roots | 20-90 | 100 |
| Bacteria | 1-2 | 2-4 |
| Actinomycetes | 0-2 | 0-4 |
| Fungi | 2-5 | 4-10 |
| Protozoa | 0-0.5 | 0-1 |
| Nematodes | 0-0.2 | 0-1/2 |
| Earthworms | 0-2.5 | 0-5 |
| Other soil animals | 0-0.5 | 0-1 |
| Viruses | Negligible | |

References

Jeffery, S., Gardi, C., Jones, A., Montanarella, L., Marmo, L., Miko, L., Ritz, K., Peres, G., Römbke, J., van der Putten, W.H. (Eds.) (2010). European Atlas of Soil Biodiversity. Publications Office of the European Union, Luxembourg, p. 128.

Killham, 1994). Soil Ecology. Cambridge UP.

Nielsen, B.O. & Nielsen, L.B. (2009). Emergence of soil nematocera (Diptera: Nematocera) in a beech stand. Entomologiske Meddelelser 77: 117-135. (in Danish with abstract in English).

Turbé et al. (2010). Soil biodiversity: functions, threats and tools for policy makers. Bio Intelligence Service, IRD and NIOO. Report for European Commission (DG Environment).

Acknowledgement

EcoFINDERS FP7 project; EU Commission grant agreement number 264465. All images are courtesy of the European Atlas of Soil Biodiversity.